


# Percolation – Summary of L1

**Aim:** Study connections between macroscopic quantities and the underlying microscopic world in the simplest not exactly solved model displaying a phase transition.

**Objective:** Gain qualitative and quantitative understanding of the phenomenon of phase transition and associated concepts such as scalefree behaviour, scaling theory, and universality.



Each site in a lattice is occupied randomly and independently with **occupation probability**  $p$ ,  $0 \leq p \leq 1$ . A **cluster** is a group of nearest-neighbour occupied sites. The **size  $s$  of a cluster** is the number of sites in the cluster. The **critical occupation probability**  $p_c$  is the occupation probability  $p$  at which an infinite cluster appears for the first time in an infinite lattice  $L = \infty$ .

## Quantities of interest

- Onset of percolation – critical occupation probability,  $p_c$ .
- Probability that a site belongs to the infinite cluster,  $P_\infty(p)$ .
- Geometry of the infinite cluster and the finite clusters.

Excluding the infinite cluster:

- Average cluster size,  $\chi(p)$ .
- Typical size of the largest cluster,  $s_\xi(p)$ .
- Typical radius of the largest cluster,  $\xi(p)$ .

## Percolation – Summary of L2

Introduced the cluster number density  $n(s, p)$  as the number of  $s$ -clusters per lattice site implying that the probability for a site to belong to any finite cluster is  $\sum_{s=1}^{\infty} sn(s, p)$ . The probability that a site belongs to the infinite cluster

$$P_{\infty}(p) = \begin{cases} 0 & \text{for } p < p_c \\ \text{nonzero} & \text{for } p > p_c, \end{cases}$$

such that

$$P_{\infty}(p) + \sum_{s=1}^{\infty} sn(s, p) = p \quad \text{valid for all } p.$$

The average cluster size

$$\chi(p) = \frac{\sum_{s=1}^{\infty} s^2 n(s, p)}{\sum_{s=1}^{\infty} sn(s, p)}.$$

Percolation in  $d = 1$  has onset of percolation at  $p_c = 1$  and

$$n(s, p) = (1 - p)^2 p^s = (1 - p)^2 \exp(-s/s_{\xi})$$

where the characteristic cluster size (typical largest cluster)


$$s_{\xi}(p) = -\frac{1}{\ln p} \propto (p_c - p)^{-1} \quad \text{for } p \rightarrow p_c^-$$

and the average cluster size

$$\chi(p) = \frac{1 + p}{1 - p} \propto (p_c - p)^{-1} \quad \text{for } p \rightarrow p_c^-.$$

The quantities  $s_{\xi}(p)$  and  $\chi(p)$  diverge as a power laws in terms of  $(p_c - p)$ , the distance of  $p$  away from  $p_c$ .

# Percolation – Summary of L3



Considered the Bethe lattice with coordination number  $z$ . Lattice contains no loops.

Critical occupation probability  $p_c = \frac{1}{z-1}$ .

The average cluster size

$$\chi(p) = \frac{p_c(1+p)}{p_c - p} \propto (p_c - p)^{-1} \quad \text{for } p \rightarrow p_c^-.$$

The probability that a site belongs to the infinite cluster

$$P_\infty(p) = \begin{cases} 0 & \text{for } p < p_c \\ p \left[ 1 - \left( \frac{1-p}{p} \right)^3 \right] & \text{for } p > p_c \end{cases}$$

$$= \begin{cases} 0 & \text{for } p < p_c \\ 6(p - p_c) & \text{for } p \rightarrow p_c^+ \end{cases}$$

Picks up abruptly at  $p = p_c$  signaling the onset of percolation.

# Percolation – Summary of L4

Introducing

$t$  = perimeter = number of empty neighbours of a cluster

$g(s, t)$  = number of different clusters with size  $s$  and perimeter  $t$ .

the general form of the cluster number density

$$n(s, p) = \sum_{t=1}^{\infty} g(s, t) (1 - p)^t p^s.$$

In a Bethe lattice  $t = s(z - 2) + 2$  and

$$\begin{aligned} n(s, p) &\propto n(s, p_c) \exp(-s/s_\xi) \\ &\propto s^{-\tau} \exp(-s/s_\xi) \end{aligned}$$

where the characteristic cluster size

$$s_\xi \propto |p_c - p|^{-2} \quad \text{for } p \rightarrow p_c.$$

Note that

$$n(s, p) = \begin{cases} s^{-\tau} & \text{for } s \ll s_\xi \\ \text{decays rapidly} & \text{for } s \gg s_\xi. \end{cases}$$

Using the ansatz above, the average cluster size

$$\begin{aligned} \chi(p) &\propto \int_1^\infty s^{2-\tau} \exp(-s/s_\xi) ds \\ &\propto s_\xi^{3-\tau} \quad \text{for } p \rightarrow p_c \\ &\propto |p_c - p|^{-1} \quad \text{for } p \rightarrow p_c \end{aligned}$$

implying that the cluster number density exponent  $\tau = 5/2$  in the Bethe lattice.

# Percolation – Summary of L5

We introduced the critical exponents characterising the percolation phase transition at  $p = p_c$ .

The exponent  $\beta$  characterises the abrupt pick up of the order parameter

$$P_{\infty}(p) = \begin{cases} 0 & \text{for } p < p_c \\ (p - p_c)^{\beta} & \text{for } p \rightarrow p_c^+. \end{cases}$$

The exponent  $\gamma$  characterises how the average cluster size diverges

$$\chi(p) \propto |p - p_c|^{-\gamma} \quad \text{for } p \rightarrow p_c.$$

The exponent  $\sigma$  characterises how the characteristic cluster size (typical size of largest cluster) diverges

$$s_{\xi}(p) \propto |p - p_c|^{-1/\sigma} \quad \text{for } p \rightarrow p_c.$$

The exponent  $\tau$  characterises the cluster number density. Explicitly, for the Bethe lattice

$$\begin{aligned} n(s, p) &= s^{-\tau} \exp(-s/s_{\xi}) && \text{for } s \gg 1 \\ &= \begin{cases} s^{-\tau} & \text{for } 1 \ll s \ll s_{\xi} \\ \text{decays rapidly} & \text{for } s \gg s_{\xi}. \end{cases} \end{aligned}$$

Critical exponents depend only on dimensionality, not the details of the lattice.

Exponent	$d = 1$	$d = 2$	$d = 3$	$d = 4$	$d = 5$	$d = 6$	<b>Bethe</b>
$\beta$	<b>0</b>	5/36	0.41	0.64	0.84	1	<b>1</b>
$\gamma$	<b>1</b>	43/18	1.80	1.44	1.18	1	<b>1</b>
$\sigma$	<b>1</b>	36/91	0.445	0.48	0.49	1/2	<b>1/2</b>
$\tau$	<b>2</b>	<b>187/91</b>	2.189	2.31	2.41	5/2	<b>5/2</b>

# Percolation – Summary of L6

The general scaling ansatz

$$n(s, p) \propto s^{-\tau} \mathcal{G}(s/s_\xi) \quad \text{for } s \gg 1$$
$$s_\xi \propto |p - p_c|^{-\frac{1}{\sigma}} \quad \text{for } p \rightarrow p_c.$$

- E.g.  $\mathcal{G}_{1d}(x) = x^2 \exp(-x)$  and  $\mathcal{G}_{\text{Bethe}}(x) = \exp(-x)$ .  
In general

$$\mathcal{G}(s/s_\xi) = \begin{cases} \mathcal{G}(0) + \mathcal{G}'(0)s/s_\xi + \frac{1}{2}\mathcal{G}''(0)(s/s_\xi)^2 + \dots & \text{for } s \ll s_\xi \\ \text{decays rapidly} & \text{for } s \gg s_\xi. \end{cases}$$

- At  $p = p_c$ ,  $s_\xi = \infty$  so the argument of  $\mathcal{G}$  is zero. Therefore

$$n(s, p_c) \propto s^{-\tau} \mathcal{G}(0) = \text{power-law decay} = \text{scale invariance}.$$

except in  $d = 1$  where  $\mathcal{G}(0) = 0$ .

- The critical occupation probability  $p_c$  depend on lattice details.  
**Non-universal quantity.**
- Critical exponents  $\tau, \sigma$  and the scaling function  $\mathcal{G}$  are independent of lattice details and depend only on dimensionality.  
**Universal quantities.**
- Plotting  $s^\tau n(s, p)$  versus  $s/s_\xi$  **all** data fall on the graph of the universal scaling function  $\mathcal{G}$ . Concept of **data collapse**.

The scaling ansatz imply the scaling relations

$$\chi(p) \propto |p_c - p|^{-\frac{3-\tau}{\sigma}} \propto |p_c - p|^{-\gamma} \quad \text{for } p \rightarrow p_c \Rightarrow \gamma = \frac{3 - \tau}{\sigma}$$
$$P_\infty(p) \propto |p_c - p|^{\frac{\tau-2}{\sigma}} \propto (p - p_c)^\beta \quad \text{for } p \rightarrow p_c^+ \Rightarrow \beta = \frac{\tau - 2}{\sigma}$$

There are only two independent critical exponents.

# Percolation – Summary of L7

The infinite cluster at  $p = p_c$  is fractal. It looks alike on all length scales  $\ell$ . Mathematically, the concept of self-similarity is expressed by the mass of the infinite cluster in a window of size  $\ell$

$$M_\infty(p_c, \ell) \propto \ell^D \quad \text{for } \ell \gg 1.$$

To discuss geometry, we introduced the **centre of mass**,  $\mathbf{r}_{cm}$ , and the **radius of gyration**,  $R_s$ , of  $s$ -clusters, where

$$\mathbf{r}_{cm} = \frac{1}{s} \sum_{i=1}^s \mathbf{r}_i, \quad R^2(s) = \frac{1}{s} \sum_{i=1}^s |\mathbf{r}_{cm} - \mathbf{r}_i|^2, \quad R_s = \sqrt{\langle R^2(s) \rangle}.$$

The mass of large but finite cluster  $1 \ll s < \infty$  at  $p = p_c$

$$M(s, p_c, \ell) = \begin{cases} \ell^D & \text{for } \ell \ll R_s - \text{fractal} \\ R_s^D & \text{for } \ell \gg R_s - \text{non-fractal} \end{cases} \\ = \ell^D m(\ell/R_s)$$

where the crossover function

$$m(\ell/R_s) \propto \begin{cases} \text{constant} & \text{for } \ell/R_s \ll 1 \\ (\ell/R_s)^{-D} & \text{for } \ell/R_s \gg 1. \end{cases}$$

The **correlation length**,  $\xi$ , is the radius of gyration of the characteristic cluster size

$$\xi = R_{s_\xi}.$$

It is the typical radius of the largest cluster. For  $p > p_c$ , it is also the typical radius of the largest hole in the infinite cluster. Since  $s_\xi$  diverges for  $p \rightarrow p_c$ , so does the correlation length and

$$\xi \propto |p - p_c|^{-\nu} \quad \text{for } p \rightarrow p_c.$$

# Percolation – Summary of L8

The correlation length,  $\xi$ , is the typical largest cluster radius:

$$\xi \propto |p - p_c|^{-\nu} \quad \text{for } p \rightarrow p_c.$$

Typical largest cluster size at occupation probability  $p$

$$s_\xi \propto \xi^D \propto |p - p_c|^{-\nu D} \propto |p - p_c|^{-1/\sigma} \quad \text{for } p \rightarrow p_c.$$

Mass of the infinite cluster for  $p \geq p_c$ :

$$\begin{aligned} M_\infty(\xi, \ell) &= \begin{cases} \ell^D & \text{for } \ell \ll \xi \\ \xi^D (\ell/\xi)^d & \text{for } \ell \gg \xi \end{cases} \\ &= \begin{cases} \ell^D & \text{for } \ell \ll \xi - \text{fractal} \\ P_\infty(p, \ell) \ell^d = \xi^{-\beta/\nu} \ell^d & \text{for } \ell \gg \xi - \text{homogeneous} \end{cases} \end{aligned}$$

Thus we have two additional scaling relations

$$D = 1/(\sigma\nu) \quad \text{and} \quad D - d = -\beta/\nu \quad (\text{valid for } d \leq 6).$$

The mass of the infinite cluster for  $p \geq p_c$

$$M_\infty(\xi, \ell) = \ell^D m_\infty(\ell/\xi)$$

with the crossover function

$$m_\infty(\ell/\xi) = \begin{cases} \text{constant} & \text{for } \ell \ll \xi \\ (\ell/\xi)^{d-D} & \text{for } \ell \gg \xi. \end{cases}$$

Alternatively,

$$\begin{aligned} M_\infty(\xi, \ell) &= b^D M_\infty(\xi/b, \ell/b) \\ &= \begin{cases} \ell^D M_\infty(\xi/\ell, 1) & \text{for } \ell \ll \xi \\ \xi^D M_\infty(1, \ell/\xi) & \text{for } \ell \gg \xi \end{cases} \\ &= \begin{cases} \ell^D & \text{for } \ell \ll \xi - \text{fractal} \\ \xi^D (\ell/\xi)^d & \text{for } \ell \gg \xi - \text{homogeneous.} \end{cases} \end{aligned}$$

# Percolation – Summary of L9

## Finite-size scaling


Assuming  $p \neq p_c$ , then  $|p - p_c| \propto \xi^{-1/\nu}$ , implying for  $L = \infty$

$$\begin{aligned} P_\infty(p) &\propto (p - p_c)^\beta && \propto \xi^{-\beta/\nu} && \text{for } p \rightarrow p_c^+ \\ \chi(p) &\propto |p - p_c|^{-\gamma} && \propto \xi^{\gamma/\nu} && \text{for } p \rightarrow p_c \\ M_k &\propto |p - p_c|^{(\tau-k-1)/\sigma} && \propto \xi^{D(1+k-\tau)} && \text{for } p \rightarrow p_c, D = 1/(\sigma\nu) \\ n(s, p) &\propto s^{-\tau} \mathcal{G}(s/s_\xi) && \propto s^{-\tau} \mathcal{G}(s/\xi^D) && \text{for } p \rightarrow p_c, s \gg 1. \end{aligned}$$

For finite system sizes  $L$  we find

$$\begin{aligned} P_\infty(\xi, L) &\propto \begin{cases} \xi^{-\beta/\nu} & \text{for } L \gg \xi - \text{constant} \\ L^{-\beta/\nu} & \text{for } L \ll \xi - \text{decaying.} \end{cases} \\ \chi(\xi, L) &\propto \begin{cases} \xi^{\gamma/\nu} & \text{for } L \gg \xi - \text{constant} \\ L^{\gamma/\nu} & \text{for } L \ll \xi - \text{increasing.} \end{cases} \\ M_k(\xi, L) &\propto \begin{cases} \xi^{D(1+k-\tau)} & \text{for } L \gg \xi - \text{constant} \\ L^{D(1+k-\tau)} & \text{for } L \ll \xi - \text{increasing.} \end{cases} \\ n(s, \xi) &\propto \begin{cases} s^{-\tau} \mathcal{G}(s/\xi^D) & \text{for } L \gg \xi, s \gg 1 - \text{cutoff constant} \\ s^{-\tau} \mathcal{G}(s/L^D) & \text{for } L \ll \xi, s \gg 1 - \text{cutoff increasing.} \end{cases} \end{aligned}$$

At  $p = p_c$ , the correlation length  $\xi = \infty$ , i.e., ALWAYS  $L \ll \xi$ . Measure critical exponents by investigating how the quantities scale with system size at  $p = p_c$ .



## Percolation – Summary of L10+L11

The fixed point equation for the rescaling transformation  $\xi = \xi/b$  has two solutions only:  $\xi = 0, \infty$ . These are associated with the solutions to the fixed point equation in  $p$ -space,  $T_b(p) = p$ , that is,  $p = 0, 1, p_c$ , representing the trivially self-similar states of the empty and fully occupied lattice and the nontrivial self-similar state at  $p = p_c$ , respectively. The critical exponent

$$\nu = \frac{\log(b)}{\left(\frac{dT_b(p)}{dp}\right)\Big|_{p_c}} \approx \frac{\log(b)}{\left(\frac{dR_b(p)}{dp}\right)\Big|_{p^*}}$$

where the rescaling transformation  $T_b$  has been substituted with a real-space renormalisation transformation  $R_b$  incorporating coarsening with rescaling. We identify  $p_c$  with  $p^*$ , the nontrivial solution to the fixed point equation

$$R_b(p^*) = p^*.$$

Often, the real-space renormalisation transformation chosen

$$R_b(p) = \begin{cases} \text{prob. of having a spanning cluster in block} \\ \text{prob. of having a majority of sites occupied in block.} \end{cases}$$

**Real-space renormalisation transformation procedure:**

- 1.** Divide the lattice into blocks of linear size  $b$ .
- 2.** Replace all sites in a block by a single block of size  $b$  occupied with probability  $R_b(p)$  according to the real-space renormalisation transformation (coarse graining procedure).
- 3.** Rescale all length scales by the factor  $b$ .

As critical exponents are determined by the large scale behaviour, they are insensitive to lattice details. They are universal.